

Please keep this brochure with your drug and read thoroughly before use. This content is simple translation text of Japanese Drug Information, and please use for reference purpose only. The official contents shall be drug information in Japanese language.

ひとみ・すこやか

Santen

第2類医薬品

Maximum concentrations of active ingredients for unpleasant tired eyes*

サンテメディカル10®

Ophthalmologic
drug

In the age of information technology, our eyes are increasingly overused. When eyes are overused, the ability to adjust focus decreases resulting in tired eye (eye fatigue).

サンテメディカル10 is an eye drop for eye fatigue containing five active ingredients including vitamin B₁₂ and neostigmine methylsulfate, effective for eye fatigue, at maximum concentrations* to improve declined focus adjustment function by working on muscles and parasympathetic nerves responsible for adjusting focus and to facilitate tissue metabolism by supplementing nutrients to eye.

Use this product to enjoy your "healthy eye" life.

*Maximum concentrations in manufacturing licence standards for nonprescription ophthalmic drugs: Vitamin B₁₂, neostigmine methyl sulfate, taurine, potassium L-aspartate, chlorpheniramine maleate

⚠ Precautions for Use!



Please consult with your doctor or pharmacist.

1. In the following cases, you should consult with your doctor or pharmacist before use.

- (1) If you are receiving medical treatment
- (2) If you or your family has allergic predisposition
- (3) If you have a history of allergic reaction to medication
- (4) If you have the following symptom
Severe pain in eye
- (5) If you have been diagnosed with the following condition
Glaucoma

2. In the following cases, you should stop using this product immediately and consult with your doctor or pharmacist taking this brochure with you.

- (1) If the following symptoms develop after use

Site	Symptom
Skin	Rash / flare, itching
Eye	Hyperaemia, itching, swelling

- (2) If blurry eye does not improve
- (3) If symptom does not improve after 5 to 6 days of use

Dosage / Administration

Instill 2 to 3 drops at a time, 5 to 6 times daily

● Please observe the following instructions.

- (1) Excessive use of this product may cause abnormal sensitivity to light or hyperaemia.
- (2) Children should use this product under supervision of their parents.
- (3) Do not allow eyelid or eyelash to touch the tip of bottle (Drug solution may be contaminated by eye mucus and various bacteria and become turbid).
Do not use this product if it looks turbid.
- (4) Do not use this product with wearing soft contact lens.
- (5) Do not use this product other than for instillation purpose.



Indications

Tired eye, blurry eye (increase in eye mucus, etc.), hyperaemia of conjunctiva, blepharitis (soreness of eyelid), itchy eye, prevention of eye disease (after swimming, after dust and sweat entering eye, etc.), ophthalmia due to light rays such as ultraviolet rays (snow ophthalmia), discomfort while wearing hard contact lens

Ingredients / Quantities

Ingredient	Quantity	Function
Vitamin B ₁₂ (Cyanocobalamin)	0.02%	Improves tired eyes by revitalizing ciliary muscle.
Neostigmine methyl sulfate	0.005%	Improves tired eye by improving the focus adjustment mechanism.
Vitamin B ₆ (Pyridoxine hydrochloride)	0.05%	Promotes tissue metabolism in eye.
Panthenol	0.05%	Promotes tissue metabolism in eye.
Potassium L-aspartate	1.0%	Increases tissue respiration in eye
Taurine	1.0%	Promotes tissue metabolism in eye.
Chlorpheniramine maleate	0.03%	Suppresses inflammation / itching of eye by suppressing actions of histamine.
ϵ -aminocaproic acid	1.0%	Suppresses production of inflammatory substances.
Dipotassium glycyrrhizinate	0.1%	Suppresses inflammation in eye.
Tetrahydrozoline hydrochloride	0.03%	Suppresses hyperaemia of conjunctiva (white part of eye).


As additives, this product contains disodium edetate hydrate, chlorobutanol, benzalkonium chloride solution, boric acid, d-borneol, l-menthol, and pH regulator.

<Notes regarding Ingredients / Quantities>

After use of this product, you may taste sweetness in your mouth. This is due to dipotassium glycyrrhizinate, one of ingredients, flowing into your mouth through lachrymal duct, and it does not mean that there is a problem with quality.

Precautions for Storage and Handling

- (1) Seal tightly and store in a cool place avoiding direct sunlight. To maintain the quality of this product, do not leave this product in a place where the temperature becomes very high, such as inside a car, near a heater. In addition, do not use this product left in a place where the temperature becomes very high since the heat may deform the bottle resulting in leaks or the heat may have compromised the quality of this product.
- (2) Keep this product away from children.
- (3) Do not transfer this product into other bottle. (This may cause misuse or compromise the quality of product.)
- (4) Do not share this product with others.
- (5) Do not use this product after expiration date. Once opened, you should use the product promptly even before expiration date.
- (6) Depending on storage condition, white crystals of this element may be formed at the tip of bottle or inside the cap. If this occurs, wipe off with clean gauze before use.
- (7) The red color of this product is due to the color of vitamin B₁₂ (cyanocobalamin). You can wash off spilled drug solution on your clothes during instillation with water.



Package: 12mL

Manufactured and marketed by
SANTEN PHARMACEUTICAL CO., LTD.
9-19, Shimoshinjo 3-chome, Higashiyodogawa-ku,
Osaka 533-8651, Japan
<http://www.santen.com/>

For more information, please visit <http://hitomi-sukoyaka.com>